

Proposition 65 Maximum Allowable Dose Level (MADL) for Reproductive Toxicity for *m*-Dinitrobenzene

March 2003

Office of Environmental Health Hazard Assessment (OEHHA) Reproductive and Cancer Hazard Assessment Section

Summary

The maximum allowable dose level (MADL) for *m*-dinitrobenzene is **38 micrograms/day (µg/d) for the oral route of exposure**, derived from an oral reproductive toxicity study in rats conducted by Linder et al. (1986). Dermal or inhalation exposure leading to an absorbed dose of 36 µg/d should be considered the maximum allowable for Proposition 65 purposes.

Background

This report describes the derivation of maximum allowable dose levels (MADLs) for *m*-dinitrobenzene (*meta*- dinitrobenzene, or *m*-DNB, CAS No. 99-65-0). *m*-DNB is used as an intermediate in organic synthesis and in the production of dyes, explosives, industrial solvents, and pesticides (HSDB, 2002). This chemical was listed under Proposition 65 (the Safe Drinking Water and Toxic Enforcement Act of 1986) as known to the State to cause reproductive toxicity (male reproductive toxicity), effective July 1, 1990. The Proposition 65 listing of *m*-DNB was based on a formal identification by the U.S. Environmental Protection Agency (U.S. EPA) of *m*-DNB as causing reproductive toxicity (male reproductive toxicity) (U.S. EPA 1989a, 1989b). U.S. EPA is an authoritative body under Proposition 65 for identification of chemicals as causing reproductive toxicity (Title 22 California Code of Regulations Section 12306 [22 CCR 12306]).

Procedures for the development of Proposition 65 MADLs are provided in regulations (22 CCR 12801 and 12803). Exposure at a level 1,000 times greater than the MADL is expected to have no observable effect. As specified in regulations, a MADL is derived from No Observable Effect Level (NOEL) based on the most sensitive study deemed to be of sufficient quality (22 CCR 12803).

Study Selection

Relevant studies on the male reproductive toxicity of *m*-DNB have been identified through literature searches. These studies, listed in the Bibliography for this document, have been reviewed and considered for the establishment of the MADL.

The male reproductive toxicity of *m*-DNB has been clearly shown in numerous toxicological studies in laboratory animals including rats and mice. Decreased testicular weights, morphological degeneration of germ cells, decreased testicular sperm head counts or epididymal sperm counts, or reduced fertility have been observed in rats or mice following oral treatment or intraperitoneal injection with *m*-DNB. Major findings from several relevant animal studies with long-term oral treatment with *m*-DNB are briefly summarized in Table 1. No male reproductive toxicity studies using inhalation or dermal exposure were found in the literature. No human data relevant to the male reproductive toxicity of *m*-DNB were found in the literature.

Table 1. Summary of the Male Reproductive Toxicity of *m*-DNB

Study Reference	Animals	Treatment	General Toxicity	Reproductive Effects & LOEL	NOEL (adjusted)
Cody et al., 1981	Rats (species not reported), 20 rats per group	Drinking water , 0, 3, 8, 20 mg/L for 16 weeks	No effect on body weights; no clinical sign of toxicity; increased spleen weights at ≥ 8 mg/L	Decreased testicular weights and "slight to moderate decrease in spermatogenesis." LOEL = 20 mg/L (3.1 mg/kg/d)	8 mg/L (1.13 mg/kg/d as reported by the authors)
Linder et al., 1986	Male Sprague-Dawley rats, 12 animals per group	Gavage , 0, 0.75, 1.5, 3.0, 6.0 mg/kg, five d/wk for 12 weeks	No effect on body weights; no clinical sign of toxicity at ≤ 3.0 mg/kg; increased spleen weights at ≥ 1.5 mg/kg.	Decreased testicular & epididymal weights; testicular atrophy; decreased testicular sperm head counts and epididymal sperm counts; decreased fertility. LOEL= 1.5 mg/kg (1.07 mg/kg/d)	0.75 mg/kg (0.54 mg/kg/d)
Irimura et al., 2000	Male Sprague-Dawley rats, 4-5 animals per group	Gavage , 0, 25, 50 mg/kg/d for four weeks or 0, 25, 50, 75 for two weeks	No sign of general toxicity.	Decreased testicular weights; testicular atrophy and decreased number of sperms in the ducts of the epididymides. LOEL= 50 mg/kg/d	25 mg/kg/d

Note: Adjusted NOELs: NOELs reported in the original reports were adjusted to mg/kg/d following the methods described in "MADL Calculation" section of this document.

The pertinent regulation specifies that the NOEL is to be based on the most sensitive study deemed to be of sufficient quality (22 CCR 12803(a)(4)). The study that meets this criterion is that by Linder et al. (1986). In this study, groups (12 animals/group) of 29-day-old male Sprague-Dawley rats were treated by gavage with *m*-DNB at doses of 0, 0.75, 1.5, 3.0, or 6.0 mg/kg, five days per week for up to twelve weeks. After ten weeks of treatment, each male was mated with two virgin females for a maximum of one week for evaluation of the effect of *m*-DNB on male fertility. The authors stated that no clinical signs of toxicity or effects on growth rate were observed in rats treated with ≤ 3.0 mg/kg *m*-DNB. The spleen weights were increased in rats treated with 1.5 or 3.0 mg/kg *m*-DNB. Due to severe toxicity including mortality observed in animals treated with 6.0

mg/kg, treatment in this group was terminated in Week 10 (after four days of breeding). Main quantitative data on the male reproductive effects of *m*-DNB as reported by the authors are summarized in Table 2. In addition, the authors observed atrophy and incomplete spermatogenesis in seminiferous tubules in animals treated with 3.0-mg/kg of *m*-DNB. One of 12 animals in the 1.5-mg/kg group exhibited decreased spermatogenesis. The authors stated that spermatogenesis in all animals dosed with 0.75-mg/kg of *m*-DNB appeared to be normal. Since no obvious effects on sperm parameters or testicular morphology were observed in rats treated with 0.75 mg/kg *m*-DNB, 0.75 mg/kg was determined to be the NOEL. This dose is equivalent to 0.54 mg/kg/d after adjusting from the dosing regimen in the study of five days per week to seven days per week.

Table 2. Major findings on the male reproductive toxicity of m-DNB from the study in rats by Linder et al. (1986)

Endpoints		Control	0.75 mg/kg	1.5 mg/kg	3.0 mg/kg
Implantation sites (No./litter)		15.6±0.4	14.5±0.7	14.1±0.6	0
Resorptions (%)		5.7±1.9	11.6±2.3	9.8±2.4	No data
No. of live pups per litter	Male	7.2±0.5	5.5±0.4*	6.6±0.6	No data
	Female	7.5±0.5	7.1±0.6	6.1±0.3	No data
	Total	14.7±0.5	12.7±0.7*	12.8±0.5*	No data
Testicular weights (g)		1.68±0.05	1.61±0.03	1.61±0.06	0.68±0.03*
Epididymal weights (g)		0.54±0.01	0.52±0.01	0.52±0.02	0.30±0.01*
Sperm parameters	No. (10 ⁶) per mg epididymal fluid	2.44±0.06	2.44±0.08	2.47±0.09	No data
	Sperm head (10 ⁶) per gram testis	147±3	141±6	115±9*	7±1*
	Motility (%)	63±2	60±3	61±4	No data
	Normal morphology (%)	95.4±2.2	97.6±0.7	96.7±0.9	15.1±13.0*

Note: *: Compared to control, p<0.05.

MADL Calculation

The NOEL is the highest dose level which results in no observable reproductive effect, expressed in milligrams of chemical per kilogram of bodyweight per day (22 CCR 12803(a)(1)). The NOEL is converted to a milligram per day dose level by multiplying the assumed human body weight by the NOEL (22 CCR 12803(b)). For male reproductive toxicity, the assumed body weight of man is 70 kg (22 CCR 12803(b)).

For the oral route of exposure, the following calculations were performed to derive the MADL for *m*-DNB, based on a NOEL of 0.75 mg/kg found in the rat study by Linder et al. (1986):

Adjusting of NOEL (0.75 mg/kg) from five days/week to seven days/week:
 $0.75 \text{ mg/kg} \times (5 \text{ days} \div 7 \text{ days}) = 0.54 \text{ mg/kg/d}$

Calculation of NOEL for a 70 kg man:
 $0.54 \text{ mg/kg/d} \times 70 \text{ kg/man} = 37.8 \text{ mg/d}$

The MADL is derived by dividing the NOEL by one thousand (1,000) to arrive at the maximum allowable dose level (22 CCR 12801(b)(1)). Thus, the adjusted NOEL is divided by 1,000 to obtain the MADL.

MADL_{oral} = $37.8 \text{ mg/d} \div 1000 = 37.8 \text{ } \mu\text{g/d}$ or **38 $\mu\text{g/d}$** after rounding.

Approximately 80-96% of *m*-DNB was absorbed when administered orally in rats (Nystrom & Rickert, 1987; McEuen et al., 1995). Using the highest reported absorption value for oral exposure, 38 $\mu\text{g/d}$ is approximately equivalent to an absorbed dose of 36 $\mu\text{g/d}$ ($38 \text{ } \mu\text{g/d} \times 96\% = 36 \text{ } \mu\text{g/d}$). No MADLs can currently be provided for dermal or inhalation exposures. However, dermal or inhalation exposure leading to an absorbed dose of 36 $\mu\text{g/d}$ should be the maximum allowable dose level for purposes of Proposition 65.

References

Cody TE, Witherup S, Hastings L, Stemmer K, Christian RT (1981). 1,3-dinitrobenzene: toxic effects in vivo and in vitro. *J Toxicol Environ Health* **7**, 829-47.

HSDB (2002). Hazardous Substances Data Bank: 1,3-Dinitrobenzene. National Library of Medicine, Bethesda, MD.

Irimura K, Yamaguchi M, Morinaga H, Sugimoto S, Kondou Y, Koida M (2000). Collaborative work to evaluate toxicity on male reproductive organs by repeated dose studies in rats. Detection of 1,3-dinitrobenzene-induced histopathological changes in testes and epididymides of rats with 2-week daily repeated dosing. *J Toxicol Sci* **25 Spec No**, 251-8.

Linder RE, Hess RA, Strader LF (1986). Testicular toxicity and infertility in male rats treated with 1,3-dinitrobenzene. *J Toxicol Environ Health* **19**, 477-89.

McEuen SF, Jacobson CF, Brown CD, Miller MG (1995). Metabolism and testicular toxicity of 1,3-dinitrobenzene in the rat: effect of route of administration. *Fundam Appl Toxicol* **28**, 94-9.

Nystrom DD, Rickert DE (1987). Metabolism and excretion of dinitrobenzenes by male Fischer-344 rats. *Drug Metab Dispos* **15**, 821-5.

U.S. EPA (1989a). U.S. Environmental Protection Agency 40 CFR Part 372, Community Right-to-know Release Reporting; Addition of Certain Chemicals; Proposed Rule, *Federal Register* **54**:16138-16142.

U.S. EPA (1989b). U.S. Environmental Protection Agency 40 CFR Part 372, Community Right-to-know Release Reporting; Addition of Certain Chemicals; Final Rule, *Federal Register* **54**:49948-49953.

Bibliography

Relevant Studies on the Male Reproductive Toxicity of *m*-DNB: The studies listed below were reviewed and considered in the establishment of the MADL.

Berger T, Miller MG, Horner CM (2000). In vitro fertilization after in vivo treatment of rats with three reproductive toxicants. *Reprod Toxicol* **14**, 45-53.

Bjorge C, Brunborg G, Wiger R, Holme JA, Scholz T, Dybing E, Soderlund EJ (1996). A comparative study of chemically induced DNA damage in isolated human and rat testicular cells. *Reprod Toxicol* **10**, 509-19.

Blackburn DM, Gray AJ, Lloyd SC, Sheard CM, Foster PM (1988). A comparison of the effects of the three isomers of dinitrobenzene on the testis in the rat. *Toxicol Appl Pharmacol* **92**, 54-64.

Brown CD, Forman CL, McEuen SF, Miller MG (1994). Metabolism and testicular toxicity of 1,3-dinitrobenzene in rats of different ages. *Fundam Appl Toxicol* **23**, 439-46.

Brown CD, Jacobson CF, Miller MG (1997). Metabolism and testicular toxicity of 1,3-dinitrobenzene in the rat: evaluation of the stage-synchrony model. *Reprod Toxicol* **11**, 57-67.

Cave DA, Foster PM (1990). Modulation of *m*-dinitrobenzene and *m*-nitrosonitrobenzene toxicity in rat Sertoli-germ cell cocultures. *Fundam Appl Toxicol* **14**, 199-207.

Cody TE, Witherup S, Hastings L, Stemmer K, Christian RT (1981). 1,3-dinitrobenzene: toxic effects in vivo and in vitro. *J Toxicol Environ Health* **7**, 829-47.

Davis RO, Gravance CG, Thal DM, Miller MG (1994). Automated analysis of toxicant-induced changes in rat sperm head morphometry. *Reprod Toxicol* **8**, 521-9.

Ellis MK, Foster PM (1992). The metabolism of 1,3-dinitrobenzene by rat testicular subcellular fractions. *Toxicol Lett* **62**, 201-8.

Evenson DP, Janca FC, Baer RK, Jost LK, Karabinus DS (1989a). Effect of 1,3-dinitrobenzene on prepubertal, pubertal, and adult mouse spermatogenesis. *J Toxicol Environ Health* **28**, 67-80.

Evenson DP, Janca FC, Jost LK, Baer RK, Karabinus DS (1989b). Flow cytometric analysis of effects of 1,3-dinitrobenzene on rat spermatogenesis. *J Toxicol Environ Health* **28**, 81-98.

Foster PM (1989). *m*-Dinitrobenzene: studies on its toxicity to the testicular Sertoli cell. *Arch Toxicol Suppl* **13**, 3-17.

Hess RA, Linder RE, Strader LF, Perreault SD (1988). Acute effects and long-term sequelae of 1,3-dinitrobenzene on male reproduction in the rat. II. Quantitative and qualitative histopathology of the testis. *J Androl* **9**, 327-42.

Holloway AJ, Moore HD, Foster PM (1990). The use of in vitro fertilization to detect reductions in the fertility of male rats exposed to 1,3-dinitrobenzene. *Fundam Appl Toxicol* **14**, 113-22.

Irimura K, Yamaguchi M, Morinaga H, Sugimoto S, Kondou Y, Koida M (2000). Collaborative work to evaluate toxicity on male reproductive organs by repeated dose studies in rats. Detection of 1,3-dinitrobenzene-induced histopathological changes in testes and epididymides of rats with 2-week daily repeated dosing. *J Toxicol Sci* **25 Spec No**, 251-8.

Jacobson CF, Miller MG (1997). 1,3-Dinitrobenzene metabolism and toxicity in seminiferous tubules isolated from rats of different ages. *Toxicology* **123**, 15-26.

Jacobson CF, Miller MG (1998). Species difference in 1,3-dinitrobenzene testicular toxicity: in vitro correlation with glutathione status. *Reprod Toxicol* **12**, 49-56.

Linder RE, Hess RA, Perreault SD, Strader LF, Barbee RR (1988). Acute effects and long-term sequelae of 1,3-dinitrobenzene on male reproduction in the rat. I. Sperm quality, quantity, and fertilizing ability. *J Androl* **9**, 317-26.

Linder RE, Hess RA, Strader LF (1986). Testicular toxicity and infertility in male rats treated with 1,3-dinitrobenzene. *J Toxicol Environ Health* **19**, 477-89.

Linder RE, Strader LF, Barbee RR, Rehnberg GL, Perreault SD (1990). Reproductive toxicity of a single dose of 1,3-dinitrobenzene in two ages of young adult male rats. *Fundam Appl Toxicol* **14**, 284-98.

Lloyd SC, Foster PM (1987). 1,3-Dinitrobenzene: toxicity and metabolism in rat testicular cell cultures. *Arch Toxicol Suppl* **11**, 281-4.

McEuen SF, Jacobson CF, Brown CD, Miller MG (1995). Metabolism and testicular toxicity of 1,3-dinitrobenzene in the rat: effect of route of administration. *Fundam Appl Toxicol* **28**, 94-9.

McEuen SF, Miller MG (1991). Metabolism and pharmacokinetics of 1,3-dinitrobenzene in the rat and the hamster. *Drug Metab Dispos* **19**, 661-6.

McLaren TT, Foster PM, Sharpe RM (1993). Identification of stage-specific changes in protein secretion by isolated seminiferous tubules from the rat following exposure to either m-dinitrobenzene or nitrobenzene. *Fundam Appl Toxicol* **21**, 384-92.

Miller MG, McEuen SF, Nasiri M, Eck DL, Kurth MJ (1991). Application of ELISA

techniques to metabolic disposition studies for 1,3-dinitrobenzene: comparison with HPLC and radiochemical methods. *Chem Res Toxicol* **4**, 324-9.

Nystrom DD, Rickert DE (1987). Metabolism and excretion of dinitrobenzenes by male Fischer-344 rats. *Drug Metab Dispos* **15**, 821-5.

Obasaju MF, Katz DF, Miller MG (1991). Species differences in susceptibility to 1,3-dinitrobenzene-induced testicular toxicity and methemoglobinemia. *Fundam Appl Toxicol* **16**, 257-66.

Reader SC, Shingles C, Stonard MD (1991). Acute testicular toxicity of 1,3-dinitrobenzene and ethylene glycol monomethyl ether in the rat: evaluation of biochemical effect markers and hormonal responses. *Fundam Appl Toxicol* **16**, 61-70.

Rehnberg GL, Linder RE, Goldman JM, Hein JF, McElroy WK, Cooper RL (1988). Changes in testicular and serum hormone concentrations in the male rat following treatment with m-dinitrobenzene. *Toxicol Appl Pharmacol* **95**, 255-64.

Strandgaard C, Miller MG (1998). Germ cell apoptosis in rat testis after administration of 1,3-dinitrobenzene. *Reprod Toxicol* **12**, 97-103.

Williams J, Foster PM (1989). The effects of 1,3-dinitrobenzene and mono-(2-ethylhexyl)phthalate on hormonally stimulated lactate and pyruvate production by rat Sertoli cell cultures. *Toxicol Lett* **47**, 249-57.