



Health Advisory and Guidelines for Eating Fish from Stevens Creek Reservoir (Santa Clara County)

June 2021



Fish, Ecotoxicology, and Water Section
Pesticide and Environmental Toxicology Branch
Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

LIST OF CONTRIBUTORS

Office of Environmental Health Hazard Assessment

Project Leads

Shannon R. Murphy, Ph.D.
Lori Chumney, M.S.

Primary Reviewers

Susan A. Klasing, Ph.D., Section Chief
Wesley Smith, Ph.D.
Huyen Tran Pham, M.P.H.

Final Reviewers

David Ting, Ph.D., Branch Chief
David Siegel, Ph.D., Assistant to the Deputy Director
Vincent Cogliano, Ph.D., Deputy Director for Scientific Programs
Sam Delson, M.A., M.P.A., Deputy Director for External and Legislative Affairs

Director

Lauren Zeise, Ph.D.

ACKNOWLEDGMENTS

Developing fish consumption advisories depends on sampling and analysis of fish. The Office of Environmental Health Hazard Assessment acknowledges the contribution of information from the following entities: the State Water Resources Control Board, the California Department of Fish and Wildlife and its analytical resources, the Moss Landing Marine Laboratories and the Water Pollution Control Laboratory. Data were obtained from the [California Environmental Data Exchange Network](http://ceden.waterboards.ca.gov/AdvancedQueryTool) (<http://ceden.waterboards.ca.gov/AdvancedQueryTool>). The map was created using ArcMap (10.5) from Environmental Systems Resource Institute (ESRI, Redlands, California).

For further information, contact:

Pesticide and Environmental Toxicology Branch
Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

1001 I Street, P.O. Box 4010
Sacramento, CA 95812-4010
Telephone: (916) 324-7572
Email address: fish@oehha.ca.gov

1515 Clay Street, 16th Floor
Oakland, California 94612
Telephone: (510) 622-3170

LIST OF ACRONYMS AND ABBREVIATIONS

ATL	Advisory Tissue Level
CDFW	California Department of Fish and Wildlife
DDT(s)	dichlorodiphenyltrichloroethane (DDT) and its metabolites dichlorodiphenyldichloroethane (DDD) and dichlorodiphenyldichloroethylene (DDE)
DHA	docosahexaenoic acid
EPA	eicosapentaenoic acid
FDA	Food and Drug Administration
Hg	mercury
MDL	method detection limit
MLML	Moss Landing Marine Laboratories
mm	millimeters
OEHHA	Office of Environmental Health Hazard Assessment
PBDEs	polybrominated diphenyl ethers
PCBs	polychlorinated biphenyls
ppb	parts per billion
RL	reporting limit
RWB2	Regional Water Board 2 (San Francisco Bay)
SCVWD	Santa Clara Valley Water District
Se	selenium
SWAMP	Surface Water Ambient Monitoring Program
SWRCB	State Water Resources Control Board
TMDL	Total Maximum Daily Load
TSMP	Toxic Substances Monitoring Program
USDA	United States Department of Agriculture
USDHHS	United States Department of Health and Human Services
US EPA	United States Environmental Protection Agency

PREFACE

The Office of Environmental Health Hazard Assessment (OEHHA), a department in the California Environmental Protection Agency, is responsible for evaluating potential public health risks from chemical contamination of sport fish. This includes issuing fish consumption advisories, when appropriate, for the State of California. OEHHA's authorities to conduct these activities are based on mandates in the:

- California Health and Safety Code
 - Section 59009, to protect public health
 - Section 59011, to advise local health authorities
- California Water Code
 - Section 13177.5, to issue health advisories

The health advisories are published in the California Department of Fish and Wildlife Sport Fishing Regulations in the section on public health advisories.

This report presents guidelines for eating fish from Stevens Creek Reservoir in Santa Clara County. The report provides background information and a technical description of how the guidelines were developed. The resulting advice is summarized in the illustrations after the Table of Contents and the List of Figures and Tables.

TABLE OF CONTENTS

A GUIDE TO EATING FISH FROM STEVENS CREEK RESERVOIR	5
INTRODUCTION	6
<i>Location</i>	6
<i>Approach Used</i>	7
CHEMICALS OF POTENTIAL CONCERN.....	7
DATA SOURCES.....	9
<i>Toxic Substances Monitoring Program (TSMP), 1989–2001</i>	9
<i>Contaminants in Fish from California Lakes and Reservoirs, 2007–2008 (SWAMP)</i>	9
<i>Long-Term Monitoring of Bass Lakes and Reservoirs in California, 2015 – ongoing (SWAMP)</i>	9
FISH SAMPLED FROM STEVENS CREEK RESERVOIR.....	10
CHEMICAL CONCENTRATIONS	10
<i>Mercury</i>	11
<i>PCBs, PBDEs, and Pesticides</i>	11
<i>Selenium</i>	11
DEVELOPMENT OF GUIDELINES FOR EATING FISH FROM STEVENS CREEK RESERVOIR.....	12
CONSUMPTION ADVICE FOR FISH FROM STEVENS CREEK RESERVOIR.....	15
<i>Black Bass Species (Largemouth Bass)</i>	15
<i>Common Carp</i>	15
<i>Crappie Species (Black Crappie)</i>	16
<i>Sacramento Sucker</i>	16
RECOMMENDED MAXIMUM NUMBER OF SERVINGS.....	17
REFERENCES	18
Stevens Creek Reservoir Fish Advisory	3

APPENDIX I. Advisory Tissue Levels.....21

LIST OF FIGURES AND TABLES

Figure 1. Location of Stevens Creek Reservoir 6

Table 1. Fish Samples Evaluated for the Stevens Creek Reservoir Advisory 10

Table 2. Mercury Concentrations in Fish from Stevens Creek Reservoir 12

Table 3. PCB Concentrations in Fish from Stevens Creek Reservoir 12

Table 4. Recommended Maximum Number of Servings per Week for Fish from Stevens Creek Reservoir 17

Advisory Tissue Levels for Selected Analytes..... 21



Women
(18 – 49 Years)

Children
(1 – 17 Years)



Women
(50+ Years)

Men
(18+ Years)

A GUIDE TO EATING FISH

from

STEVENS CREEK RESERVOIR

(Santa Clara County)

Eat the Good Fish
Eating fish that are low in chemicals may provide health benefits to children and adults.



Avoid the Bad Fish
Eating fish with higher levels of chemicals like mercury or PCBs may cause health problems in children and adults.



Choose the Right Fish
Chemicals may be more harmful to unborn babies and children.



Common Carp



Sacramento Sucker



Crappie Species



Black Bass Species

1 TOTAL SERVING A WEEK

2 TOTAL SERVINGS A WEEK

OR

0 DO NOT EAT

1 TOTAL SERVING A WEEK

0 DO NOT EAT

0 DO NOT EAT

Serving Size
A serving of fish is about the size and thickness of your hand. Give children smaller servings.

For Adults



For Children



California Office of Environmental Health Hazard Assessment

web www.oehha.ca.gov/fish
 email fish@oehha.ca.gov
 phone (916) 324-7572

Eat only the skinless fillet



Eat only the meat



Some chemicals are higher in the skin, fat, and guts.
Updated 06/2021

INTRODUCTION

This report updates and supersedes the previous guidelines (OEHHA, 2009) for eating fish from Stevens Creek Reservoir. This update provides new advice for Common Carp and Sacramento Sucker. Consumption advice for eating black bass species, Common Carp, crappie species, and Sacramento Sucker, is based on levels of mercury found in fish collected from Stevens Creek Reservoir.

LOCATION

Stevens Creek Reservoir is located in the foothills of the Santa Cruz Mountains about 15 miles west of San Jose, CA (Figure 1). Stevens Creek Reservoir was formed in 1935 by construction of an earthen dam on Stevens Creek¹. Swiss Creek also feeds this 3,138 acre-foot reservoir that serves to restore local aquifers and provide recreation for fishing and non-powered boating activities (SCVWD, 2017a,b and 2020). Santa Clara Valley Water District (SCVWD) manages Stevens Creek Dam, and Santa Clara County Parks² manages Stevens Creek County Park, including the reservoir.

FIGURE 1. LOCATION OF STEVENS CREEK RESERVOIR



¹ Information about Stevens Creek Reservoir is available at: <https://www.valleywater.org/your-water/local-dams-and-reservoirs>.

² Information from Santa Clara County Parks available at: <https://www.sccgov.org/sites/parks/parkfinder/Pages/StevensCreek.aspx>.

APPROACH USED

The Office of Environmental Health Hazard Assessment (OEHHA) used the results from three monitoring studies described in this report to develop the Stevens Creek Reservoir Advisory. OEHHA uses the following general process in developing consumption advice for sport fish:

- 1) Evaluation of all fish contaminant data available from a water body and selection of appropriate data that meet data quality criteria and sampling plan guidelines.
- 2) Determination of fish species for which adequate data are available to issue fish consumption advice.
- 3) Calculation of an appropriate measure of central tendency (often a weighted arithmetic mean³) and other descriptive statistics of the contaminant data, as appropriate, for a chemical of potential concern for the selected fish species.
- 4) Comparison of the chemical concentrations with the OEHHA Advisory Tissue Levels (ATLs) for each chemical of potential concern.
- 5) Development of final advice based on a thorough review of the data and best professional judgment relating to the benefits and risks of consuming a particular fish species.

The ATLs (discussed further in a subsequent section of this report) are chemical levels in fish tissue that are considered acceptable, based on chemical toxicity, for a range of consumption rates. Development of the ATLs also included consideration of health benefits associated with including fish in the diet (OEHHA, 2008). The ATLs should not be interpreted as static “bright lines,” but one component of a complex process of data evaluation and interpretation used by OEHHA in the assessment and communication of the benefits and risks of consuming sport fish.

CHEMICALS OF POTENTIAL CONCERN

Certain chemicals are of potential concern for people who eat fish because of their toxicity and their ability to accumulate in fish tissue. The majority of fish consumption advisories in California are issued because of mercury (Hg), followed by polychlorinated biphenyls (PCBs) and, in a few cases, selenium (Se), polybrominated diphenyl ethers (PBDEs), or some legacy pesticides (pesticides that are no longer used but remain in the environment).

Mercury is a natural element found in some rock and soil. Human activities, such as burning coal and the historical use of mercury to mine gold, also add mercury to the environment. If mercury enters waterways, it can be converted to a more toxic form

³ Means are an arithmetic average of individual values and/or a composites weighted by number of fish. A weighted average of composites is calculated by multiplying the chemical concentration in each composite by the number of fish in that composite for each species. Products are then summed and divided by the total number of fish in all composites for that species.

known as methylmercury – which can pass into and build up in fish. High levels of methylmercury can harm the brain, especially in fetuses and children.

PCBs are industrial chemicals previously used in electrical transformers, plastics, and lubricating oils, often as flame retardants or electrical insulators. Their use was banned in the 1970s, but they persist in the environment because they do not break down easily and can accumulate in fish. Depending on the exposure level, PCBs may cause cancer or other health effects, including neurotoxicity, in humans.

Selenium is a naturally occurring metalloid and at low doses is an essential nutrient for many important human health processes, including thyroid regulation and vitamin C metabolism. Higher doses cause selenium toxicity, which can include symptoms ranging from hair loss and gastrointestinal distress to dizziness and tremors.

Chlordanes, dichlorodiphenyltrichloroethane (DDT), dieldrin, and toxaphene are pesticides that were banned from use in 1973 (DDT), the late 1980s (chlordanes and dieldrin) and 1990 (toxaphene), but are still found in some fish in certain California water bodies. Depending on the exposure level, these chemicals may cause cancer or adverse effects on the nervous system.

PBDEs are a class of flame retardants historically used in a variety of consumer products, including furniture, textiles, automotive parts, and electronics. The use of PBDEs in new products was largely phased out by 2013 but, due to their wide usage and persistence in the environment, they are still being detected in fish tissues. PBDEs may affect hormone levels or learning and behavior in children.

Detailed discussion of the toxicity of these chemicals and references are presented in “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, Dieldrin, Methylmercury, PCBs, Selenium, and Toxaphene” (OEHHA, 2008) and “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Polybrominated Diphenyl Ethers (PBDEs)” (OEHHA, 2011).

All fish species collected from Stevens Creek Reservoir and used in advisory development were analyzed for mercury (as a measure of methylmercury). Common Carp and Largemouth Bass were analyzed for the legacy pesticides, chlordanes (cis-chlordane, trans-chlordane, cis-nonachlor, trans-nonachlor, and oxychlordane), dieldrin, DDTs (DDT and its metabolites dichlorodiphenyldichloroethane [DDD], dichlorodiphenyldichloroethylene [DDE]), and toxaphene (Largemouth Bass only). Common Carp were also analyzed for PBDEs, PCBs, and selenium. Selenium was also analyzed in Sacramento Sucker samples. Fish species that do not normally accumulate PCBs or other organic chemicals may not be analyzed for those contaminants in a particular monitoring study. Additionally, some studies do not analyze these chemicals and instead focus only on mercury.

DATA SOURCES

The guidelines for eating fish from Stevens Creek Reservoir are based on the chemicals detected in the fish collected for the three monitoring studies described below. These studies met OEHHA's data quality criteria, including adequate documentation of sample collection, fish preparation methods (e.g., skinning or filleting), chemical analyses, quality assurance, and sufficiently low detection limits. "Sample," as used in this report, refers to an individual fish or a composite of multiple fish for which contaminant data were reported. "Sampling" or "sampled" refers to the act of collecting fish for chemical analysis.

TOXIC SUBSTANCES MONITORING PROGRAM (TSMP), 1989–2001

The TSMP operated from 1976 to 2003 as a state water quality-monitoring program managed by SWRCB (SWRCB, 2007 and 2013). Its objective was to provide statewide information on the occurrence of toxic substances by monitoring water bodies with known or suspected water quality impairment. As part of this program, the California Department of Fish and Wildlife (CDFW), then known as the California Department of Fish and Game, collected Black Crappie and Largemouth Bass in 2001, and Sacramento Sucker in 1989 from Stevens Creek Reservoir. All fish samples were analyzed for mercury, Sacramento Sucker were also analyzed for selenium, and Largemouth Bass analysis included the legacy pesticides, chlordanes, DDTs, dieldrin, and toxaphene.

CONTAMINANTS IN FISH FROM CALIFORNIA LAKES AND RESERVOIRS, 2007–2008 (SWAMP)

The Surface Water Ambient Monitoring Program (SWAMP), operated by the State Water Resources Control Board (SWRCB) in cooperation with the San Francisco Bay Regional Water Quality Control Board (RWB2), monitors water quality in California's surface waters. This survey of inland water bodies was the State's largest survey of chemical contaminants in sport fish, sampling 272 lakes and reservoirs from 2007 to 2008 (SWRCB, 2010). The SWRCB used the data from this survey to characterize statewide water quality conditions. The program collected Largemouth Bass (mercury analysis only) and Common Carp from Stevens Creek Reservoir in 2007, where Common Carp were analyzed for chlordanes, DDTs, dieldrin, mercury, PBDEs, PCBs, and selenium.

LONG-TERM MONITORING OF BASS LAKES AND RESERVOIRS IN CALIFORNIA, 2015 – ONGOING (SWAMP)

This monitoring study is a multi-year effort initiated in 2015 to document long-term trends related to mercury contamination in sport fish from California lakes and reservoirs dominated by bass species (Davis et al., *in preparation*). In 2019, SWAMP collected Largemouth Bass from Stevens Creek Reservoir, which were analyzed for mercury.

FISH SAMPLED FROM STEVENS CREEK RESERVOIR

The fish sampling data used in this advisory were retrieved from the California Environmental Data Exchange Network (CEDEN), the state's repository for environmental data. Samples were excluded when the fish were not legal size to take or did not meet OEHHA's criteria for minimum "edible" size based on species size at maturity, and professional judgment (as described in OEHHA, 2005). A summary of all fish species evaluated for this advisory is shown in Table 1, including the name of the species, number of samples collected, total number of fish, project name, year sampled, and contaminants analyzed.

TABLE 1. FISH SAMPLES EVALUATED FOR THE STEVENS CREEK RESERVOIR ADVISORY

Common Name	Scientific Name	Number of Samples	Total Number of Fish	Project	Year Collected	Contaminants Analyzed ^b
Black Crappie	<i>Pomoxis nigromaculatus</i>	4	20	TSMP ^a	2001	Hg
Common Carp	<i>Cyprinus carpio</i>	2	10	SWAMP	2007	Chlordanes, DDTs, Dieldrin, Hg, PBDEs, PCBs, Se
Largemouth Bass	<i>Micropterus salmoides</i>	3	12	TSMP ^a	2001	Chlordanes, DDTs, Dieldrin, Hg, Toxaphene
		1	6	SWAMP	2007, 2019	Hg
Sacramento Sucker	<i>Catostomus occidentalis</i>	1	9	TSMP ^a	1989	Hg, Se

Samples were analyzed as skinless fillets, with the following exceptions:

^aStudy report did not specify whether skin was removed from fillets prior to tissue analysis.

^bOrganic data (chlordanes, DDTs, dieldrin, PCBs or toxaphene) generated prior to 2000 were excluded from the analysis because data that are more recent are considered more reliable due to improved analytical methods.

CHEMICAL CONCENTRATIONS

As shown in Table 1, samples were analyzed for one or more of the following: total mercury, selenium, chlordanes, DDTs, dieldrin, toxaphene, PBDEs (7 congeners), and PCBs (54–55 congeners)⁴. Among the chemicals analyzed in fish tissue samples from Stevens Creek Reservoir, only mercury was sufficiently high to impact consumption advice.

⁴ Congeners are related compounds with similar chemical forms. Of the 209 possible PBDE and PCB congeners, 6–7 and 48–54 are generally analyzed, respectively.

All fish samples were prepared as skinless fillets, except for the TSMP study where the fillet preparation method for Black Crappie and Largemouth Bass was not recorded. Samples were analyzed as individual fish or composites.

For this advisory, OEHHA used the weighted (by the number of individual fish) arithmetic mean (average) of the chemical concentrations (in wet weight) for each fish species to estimate average human exposure.

MERCURY

Samples were analyzed for total mercury, as either individual fish or composite samples, using a direct mercury analyzer (DMA) at the CDFW Moss Landing Marine Laboratories (MLML). The DMA method utilizes thermal decomposition and atomic absorption. OEHHA assumed all mercury detected was methylmercury, which is the most common form found in fish and is also the more toxic form (Bloom, 1992). Table 2 shows the averages and ranges for total length⁵, as well as mercury concentrations in each fish species. The DMA method detection limit (MDL)⁶ and the reporting limit (RL)⁷ for total mercury were reported at 3 or 12 and 9 or 12 parts per billion (ppb), respectively, depending on the study. Although mercury was detected at commonly found concentrations in the TSMP study, the MDL and RL for mercury were not reported.

PCBS, PBDES, AND PESTICIDES

Pesticides, PBDEs and PCBs were analyzed by gas chromatography at the CDFW Water Pollution Control Laboratory. For chlordanes, DDTs, PCBs, and PBDEs, each of the concentrations presented was the sum of the detected parent compound, congeners, or metabolites, where applicable. Because the MDLs or RLs were generally low (≤ 5 ppb, with a few exceptions), individual congeners or metabolites with concentrations reported as non-detects were assumed to be zero. This is a standard method of handling non-detect values for PCBs and other chemicals with multiple congeners or metabolites in a given sample when detection levels are adequate (US EPA, 2000a). Table 3 shows the averages and ranges for total length, as well as PCB concentrations in Common Carp.

SELENIUM

The CDFW MLML analyzed Common Carp collected from Stevens Creek Reservoir for selenium, as composite samples, using inductively coupled plasma-mass spectrometry (ICP-MS). The ICP-MS method uses desolvation, atomization, and ionization with ion

⁵ Total length is the maximum length of the fish, measured from the tip of the closed mouth to the tip of the pinched tail fin.

⁶ The MDL is the lowest quantity of a chemical that can be distinguished (as greater than zero) in a sample.

⁷ The RL is the lowest quantity of a chemical that can be accurately quantified in a sample.

separation based on a mass-to-charge ratio to detect the total selenium concentration in a sample. The ICP-MS MDL and RL for total selenium for the SWAMP study were 100 and 300, respectively.

Concentrations of chlordanes, dieldrin, DDTs, PBDEs, selenium, and toxaphene were lower than the corresponding ATL threshold values for daily consumption (OEHHA, 2008 and 2011). These chemicals were therefore not considered further for developing consumption advice and are not shown in this report.

TABLE 2. MERCURY CONCENTRATIONS IN FISH FROM STEVENS CREEK RESERVOIR

Species from Stevens Creek Reservoir	Number of Samples ^a	Total Number of Fish	Mean* Total Length (mm)	Range of Total Lengths** (mm)	Mercury (ppb)	
					Mean*	Range**
Black Crappie	4	20	206	201 – 210	597	557 – 616
Common Carp	2	10	604	542 – 634	306	292 – 320
Largemouth Bass	9	18	440	330 – 490	1306	568 – 1630
Sacramento Sucker	1	9	326	n/a	340	n/a

^aSamples were prepared as skinless filets, except as noted in the footnotes to Table 1.

*Means are an arithmetic average of individual values and/or a weighted average of composites.

**Range of individuals and/or range of the composites.

n/a = not applicable due to a single sample

TABLE 3. PCB CONCENTRATIONS IN FISH FROM STEVENS CREEK RESERVOIR

Species from Stevens Creek Reservoir	Number of Samples ^a	Total Number of Fish	Mean* Total Length (mm)	Range of Total Lengths** (mm)	PCBs (ppb)	
					Mean*	Range**
Common Carp	2	10	604	542 – 634	19	16 – 23

^aSamples were prepared as skinless filets.

*Means are an arithmetic weighted average of composites.

**Range of the composites.

DEVELOPMENT OF GUIDELINES FOR EATING FISH FROM STEVENS CREEK RESERVOIR

The OEHHA fish advisory process considers the health benefits of fish consumption as well as the risk from exposure to the chemical contaminants found in fish. Benefits are included in the advisory process because there is considerable evidence and scientific consensus that fish should be part of a healthy well-balanced diet. Fish contain many

nutrients that are important for general health and, in particular, help promote optimal growth and development of babies and young children, and may reduce the incidence of heart disease in adults (FDA/US EPA, 2017; American Heart Association, 2016; OEHHA, 2008; Institute of Medicine, 2007; Kris-Etherton et al., 2002). Fish are a significant source of the beneficial omega-3 fatty acids, docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) (USDA/USDHHS, 2020; Weaver et al., 2008).

The US Department of Agriculture (USDA) recommends “including at least 8 ounces of cooked seafood⁸ per week. Young children need less, depending on their age and calorie needs” (MyPlate.gov). According to the 2020–2025 Dietary Guidelines, “women who are pregnant or lactating should consume at least 8 and up to 12 ounces of a variety of seafood per week from choices that are lower in methylmercury” (USDA/USDHHS, 2020). Additionally, “based on FDA and EPA’s advice, depending on body weight, some women should choose seafood lowest in methylmercury or eat less seafood than the amounts in the Healthy U.S.-Style Dietary Pattern” (USDA/USDHHS, 2020). For more-detailed information, see USDA/USDHHS (2020) and other USDA MyPlate.gov materials. The particular fish that people eat is an important factor in determining the net beneficial effects of fish consumption. For example, studies have shown that children of mothers who ate low-mercury fish during pregnancy scored better on cognitive tests compared to children of mothers who did not eat fish or ate high-mercury fish (Oken et al., 2005 and 2008). Accordingly, because of the high mercury content of certain fish species, the US Food and Drug Administration (FDA) and the US Environmental Protection Agency recommend that women who are pregnant (or might become pregnant) or breastfeeding, and young children avoid consuming shark, swordfish, tilefish (Gulf of Mexico), bigeye tuna, marlin, orange roughy, and king mackerel (FDA/US EPA, 2017).

To address the potential health concerns associated with exposure to contaminants in sport fish, OEHHA has established ATLS for chemicals that are known to accumulate in the edible tissues of fish. ATLS consider both the toxicity of the chemical and potential benefits of eating fish. OEHHA uses the ATLS to determine the maximum number of servings per week that consumers should eat, for each species and at each location, to limit their exposure to these contaminants. Consumers can use OEHHA’s guidance when choosing which fish and how much to eat as part of an overall healthy diet.

There are two sets of ATLS for methylmercury in fish because of the age-related toxicity of this chemical (OEHHA, 2008). The fetus and children are more sensitive to the toxic effects of methylmercury. Thus, the ATLS for the sensitive population, including women who might become pregnant (typically 18 to 49 years of age) and children 1-17 years, are lower than those for women 50 years and older and men 18 years and older. The lower ATL values for the sensitive population provide additional protection to allow for normal growth and development of the brain and nervous system of unborn babies and children. Detailed discussion about the toxicity of common fish contaminants and health benefits of fish consumption, as well as derivation of the ATLS, are provided in

⁸ Seafood as used here refers to fish and shellfish from freshwater and marine environments.

“Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, Dieldrin, Methylmercury, PCBs, Selenium, and Toxaphene” (OEHHA, 2008) and “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Polybrominated Diphenyl Ethers (PBDEs)” (OEHHA, 2011). A list of the ATLS used in this report is presented in Appendix I.

For each fish species in this advisory, OEHHA compared the mean chemical concentrations detected in the fillet to the corresponding ATLS to establish the maximum number of servings per week that could be consumed (see Appendix I). For fish fillets, a serving size is considered to be 8 ounces, prior to cooking, or about the size and thickness of a hand. Children should be given smaller servings. For smaller fish species, several individual fish may be required to yield a serving.

The consumption advice for a fish species is initially based on the chemical with the lowest allowable number of servings per week. Because some chemicals, such as mercury and PCBs, are known to have similar adverse effects, additivity of toxicity is assumed in such cases and may be assessed using multiple chemical exposure methodology (US EPA, 1989 and 2000b). If two or more chemicals with similar adverse effects are present in fish tissue, multiple-chemical-exposure methodology is employed. This may result in advising fewer servings per week than would be the case for the presence of one chemical alone, in a similar concentration. The potential effect of multiple chemical exposures (mercury and PCBs) was assessed in Common Carp and did not affect advice. Advice for all species in this advisory was based solely on mercury concentrations without the need to apply the multiple-chemical method.

OEHHA recommends that individuals strive to meet the US Dietary Guidelines’ seafood consumption recommendations, while also adhering to federal and OEHHA recommendations to limit the consumption of fish with higher contaminant levels. The advice discussed in the following section represents the maximum recommended number of servings per week for different fish species. People should eat no more than the recommended number of servings for each fish species or species group. OEHHA’s consumption advice for a particular fish species can be extended to other closely related fish species⁹ known to accumulate similar levels of contaminants.

Consumption advice should not be combined. That is, if a person chooses to eat a serving of fish from the “one-serving-a-week” category, then they should not eat any other fish from any source (including commercial) until the next week. If a person chooses to eat a serving of fish from the “two-servings-per-week” category, they can combine fish species from that category, or eat one serving of fish from that category and one from a category that recommends more than two servings per week (if available), for a total of two servings in that week. Then they should not eat any other fish from any source (including commercial) until the following week.

⁹ Fish species within the same genus are most closely related, and family is the next level of relationship.

CONSUMPTION ADVICE FOR FISH FROM STEVENS CREEK RESERVOIR

OEHHA's advisory protocol requires at least nine fish of a species to be collected from a water body before an advisory can be developed for the primary contaminant of concern. This is to ensure the sample dataset is representative of the fish species population in the water body. In some cases, an exception is made to develop advice for species that are commonly caught and consumed from a given water body but where available data may be limited. Generally, this practice applies when the advice supports no consumption of that species. For Stevens Creek Reservoir, the sample size criterion was met for the following species: Black Crappie, Common Carp, Largemouth Bass, and Sacramento Sucker. There were not sufficient data to evaluate other species that may be found in this water body. For fish species found in Stevens Creek Reservoir that are not included in this advisory, OEHHA recommends following the [statewide advisory for lakes and reservoirs without site-specific advice](#).

The following advice is based solely on mercury concentrations. The sensitive population is defined as women 18 to 49 years and children 1 to 17 years, and the general population is defined as women 50 years and older and men 18 years and older.

BLACK BASS SPECIES (LARGEMOUTH BASS)

OEHHA has evaluated mercury concentrations in black bass species in many water bodies in California and has found a similar range of mercury concentrations when two or more of these species were caught from the same water body. Therefore, OEHHA extends the consumption advice for Largemouth Bass to other black bass species, including Redeye, Smallmouth, and Spotted Bass.

Based on the mean mercury concentration of 1306 ppb in Largemouth Bass, OEHHA recommends no consumption of black bass species from Stevens Creek Reservoir for both the sensitive and general populations. Although the concentration of mercury was just below the no consumption threshold for the general population (1310 ppb), OEHHA determined that no consumption advice was appropriate for the general population based on evaluation of the range of mercury concentrations in Largemouth Bass. Out of a total of 9 samples (comprised of 18 fish), five samples (comprised of 13 fish) exceeded the no consumption threshold for mercury (1310 ppb). PCBs were not analyzed in Largemouth Bass from Stevens Creek Reservoir.

COMMON CARP

The mean mercury and PCB concentrations in Common Carp from Stevens Creek Reservoir were 306 and 19 ppb, respectively. OEHHA recommends a maximum of one serving a week of Common Carp for the sensitive population, and a maximum of two servings a week for the general population, based on mercury. PCBs did not impact advice.

CRAPPIE SPECIES (BLACK CRAPPIE)

The mean mercury concentration in Black Crappie from Stevens Creek Reservoir was 597 ppb. OEHHA recommends no consumption of Black Crappie for the sensitive population, and a maximum of one serving a week for the general population, based on mercury. PCBs were not analyzed in Black Crappie from Stevens Creek Reservoir.

OEHHA has evaluated mercury concentrations in crappie species in many water bodies in California and has found a similar range of mercury concentrations when two of these species were caught from the same water body. Therefore, OEHHA extends the consumption advice for Black Crappie to White Crappie.

SACRAMENTO SUCKER

The mean mercury concentration in Sacramento Sucker from Stevens Creek Reservoir was 340 ppb. OEHHA recommends a maximum of one serving a week of Sacramento Sucker for the sensitive population, and a maximum of two servings a week for the general population. PCBs were not analyzed in Sacramento Sucker from Stevens Creek Reservoir.

RECOMMENDED MAXIMUM NUMBER OF SERVINGS

The recommended maximum numbers of servings per week for fish from Stevens Creek Reservoir are shown in Table 4.

TABLE 4. RECOMMENDED MAXIMUM NUMBER OF SERVINGS PER WEEK FOR FISH FROM STEVENS CREEK RESERVOIR

Fish Species	Women 18–49 years and Children 1–17 years	Women 50 years and older and Men 18 years and older
Black Bass Species	0	0
Common Carp	1	2
Crappie Species	0	1
Sacramento Sucker	1	2

REFERENCES

- American Heart Association. 2016. Fish and Omega-3 Fatty Acids. Online at: http://www.heart.org/HEARTORG/HealthyLiving/HealthyEating/HealthyDietGoals/Fish-and-Omega-3-Fatty-Acids_UCM_303248_Article.jsp#.Wl57BnlG2Uk.
- Bloom, N.S. 1992. On the chemical form of mercury in edible fish and marine invertebrate tissue. *Can. J. Fish. Aquat. Sci.* 49(5):1010-1017.
- Davis, J.A., A. Bonnema, B. Jakl, G. Ichikawa, and W.A. Heim. Long-Term Monitoring of Bass Lakes and Reservoirs in California: 2019 Data Report. A Report of the Surface Water Ambient Monitoring Program (SWAMP). California State Water Resources Control Board, Sacramento, CA. *Manuscript in Preparation*.
- FDA/US EPA. 2017. Eating Fish: What pregnant women and parents should know. Advice by FDA and US EPA/January, 2017. Online at: <https://www.fda.gov/food/consumers/advice-about-eating-fish>
- Institute of Medicine. 2007. Seafood choices, balancing benefits and risks. Committee on Nutrient Relationships in Seafood: Selections to Balance Benefits and Risks. Institute of Medicine, Food and Nutrition Board. The National Academies Press, Washington, D.C.
- Kris-Etherton, P.M., W.S. Harris, and L.J. Appel. 2002. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circ.* 106:2747-2757.
- OEHHA. 2005. General Protocol for Sport Fish Sampling and Analysis. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <http://oehha.ca.gov/media/downloads/fish/document/fishsamplingprotocol2005.pdf>.
- OEHHA. 2008. Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, Dieldrin, Methylmercury, PCBs, Selenium, and Toxaphene. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <http://oehha.ca.gov/media/downloads/fish/report/atlmhgandothers2008c.pdf>.
- OEHHA. 2009. 2009 Update of California Sport Fish Advisories. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: https://oehha.ca.gov/media/downloads/advisories/discadvupdates031309_9.pdf
- OEHHA. 2011. Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Polybrominated diphenyl ethers (PBDEs). Office of Environmental Health Hazard Assessment, California Environmental

Protection Agency, Sacramento, California. Online at:
<http://oehha.ca.gov/media/downloads/fish/report/pbdes052311.pdf>.

Oken, E., R.O. Wright, K.P. Kleinman, D. Bellinger, C.J. Amarasiriwardena, H. Hu, J.W. Rich-Edwards, and M.W. Gillman. 2005. Maternal fish consumption, hair mercury, and infant cognition in a U.S. cohort. *Environ. Health Perspect.* 113(10):1376-1380.

Oken, E., J.S. Radesky, R.O. Wright, D. Bellinger, C.J. Amarasiriwardena, K.P. Kleinman, H. Hu, J.W. Rich-Edwards, and M.W. Gillman. 2008. Maternal fish intake during pregnancy, blood mercury levels, and infant cognition at age 3 years in a U.S. cohort. *Am. J. Epidemiol.* 167(10):1171-1181.

SCVWD. 2017a. Guadalupe River Watershed Mercury TMDL: 2016-2017 Progress Report on Methylmercury Production and Control Measures. Santa Clara Valley Water District. Online at:
https://www.waterboards.ca.gov/rwqcb2/water_issues/programs/TMDLs/guadalupe_river_mercury/180410-BiennialReport_20180326.pdf.

SCVWD. 2017b. 2016 and 2017 Fish Assemblage Report for the Guadalupe River Watershed Mercury Total Maximum Daily Load. Santa Clara Valley Water District. Online at:
https://www.valleywater.org/sites/default/files/B1_2016%20Reservoir%20Assemblage%20Report_FinalV2.pdf.

SCVWD. 2020. Guadalupe River Watershed TMDL 2018-2019 Progress Report on Methylmercury Production and Control Measures Supplemental Information. Santa Clara Valley Water District. Online at: <https://s3.us-west-2.amazonaws.com/assets.valleywater.org/2019%20Biennial%20Report%20Cover%20Letter%20and%20Attachments%20Final%20reduced.pdf>

SWRCB. 2007. Bioaccumulation of Pollutants in California Waters: A Review of Historic Data and Assessment of Impacts on Fishing and Aquatic Life. State Water Resources Control Board, California Environmental Protection Agency, Sacramento, California. Online at:
http://www.waterboards.ca.gov/water_issues/programs/swamp/docs/bop/cw117_swrcb_report.pdf.

SWRCB. 2010. Contaminants in Fish from California Lakes and Reservoirs, 2007-2008: Summary Report on a Two-Year Screening Survey. State Water Resources Control Board, California Environmental Protection Agency, Sacramento, California. Online at:
http://www.waterboards.ca.gov/water_issues/programs/swamp/docs/lakes_study/lake_survey_yr2_no_app.pdf.

SWRCB. 2013. State Mussel Watch (SMW) Program/Toxic Substances Monitoring (TSM) Program. State Water Resources Control Board, California Environmental

Protection Agency, Sacramento, California. Online at:
http://www.waterboards.ca.gov/water_issues/programs/swamp/mussel_watch.shtml.

USDA/USDHHS. 2020. Dietary Guidelines for Americans, 2020-2025. 9th Edition. U.S. Department of Health and Human Services and U.S. Department of Agriculture. Online at: <https://www.dietaryguidelines.gov/>.

US EPA. 1989. Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual (Part A) Interim Final. EPA/5401-89/002, December 1989. Office of Emergency and Remedial Response, U.S. Environmental Protection Agency, Washington, D.C. Online at: <https://rais.ornl.gov/documents/HHEMA.pdf>.

US EPA. 2000a. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories: Volume 1. Fish Sampling and Analysis, 3rd Edition. EPA 823-B00-007. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

US EPA. 2000b. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories: Volume 2. Risk Assessment and Fish Consumption Limits, 3rd Edition. EPA 823-B-00-007. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

Weaver, K.L., P. Ivester, J.A. Chilton, M.D. Wilson, P. Pandey, and F.H. Chilton. 2008. The content of favorable and unfavorable polyunsaturated fatty acids found in commonly eaten fish. J. American Dietetic Assoc. 108:1178-1185.

APPENDIX I. ADVISORY TISSUE LEVELS

Advisory Tissue Levels (ATLs; OEHHA, 2008 and 2011) guide the development of advice for people eating sport fish. ATLs are levels of contaminants found in fish that correspond to the maximum numbers of recommended fish servings. OEHHA uses ATLs to provide advice to prevent consumers from being exposed to:

- More than the reference dose¹⁰ on an average daily basis for chemicals not known to cause cancer, such as methylmercury, or
- For cancer-causing chemicals, a risk level greater than one additional cancer case in a population of 10,000 people consuming fish at the given consumption rate over a lifetime. This cancer endpoint is the maximum acceptable risk level recommended by the US EPA (2000b) for fish advisories.

For each chemical, ATLs were determined for both cancer and non-cancer risk, if appropriate, for one to seven eight-ounce servings per week. The most health protective ATLs for each chemical, selected from either cancer or non-cancer based risk, are shown in the table below for zero to seven servings per week. When the guidelines for eating fish are followed, exposure to chemicals in fish would be at or below the average daily reference dose or the cancer risk probability of one in 10,000.

ADVISORY TISSUE LEVELS FOR SELECTED ANALYTES

Contaminant	Consumption Frequency Categories (8-ounce servings/week) ^a and ATLs (in ppb)							
	7	6	5	4	3	2	1	0
Chlordanes	≤ 80	>80–90	>90–110	>110–140	>140–190	>190–280	>280–560	>560
DDTs	≤ 220	>220–260	>260–310	>310–390	>390–520	>520–1,000	>1,000–2,100	>2,100
Dieldrin	≤ 7	>7–8	>8–9	>9–11	>11–15	>15–23	>23–46	>46
MeHg (Women 18–49 and children 1–17)	≤ 31	>31–36	>36–44	>44–55	>55–70	>70–150	>150–440	>440
MeHg (Women ≥ 50 and men ≥ 18)	≤ 94	>94–109	>109–130	>130–160	>160–220	>220–440	>440–1,310	>1,310
PBDEs	≤ 45	>45–52	>52–63	>63–78	>78–100	>100–210	>210–630	>630
PCBs	≤ 9	>9–10	>10–13	>13–16	>16–21	>21–42	>42–120	>120
Selenium	≤ 1000	>1,000–1200	>1,200–1,400	>1,400–1,800	>1,800–2,500	>2,500–4,900	>4,900–15,000	>15,000
Toxaphene	≤ 87	>87–100	>100–120	>120–150	>150–200	>200–300	>300–610	>610

^a Serving sizes (prior to cooking, wet weight) are based on an average 160-pound person. Individuals weighing less than 160 pounds should eat proportionately smaller amounts.

¹⁰ The reference dose is an estimate of the maximum daily exposure to a chemical likely to be without significant risk of harmful health effects over a lifetime.